
AD-A139 276 HIGH LATITUDE ELECTRODYNAMICS(U) AIR FORCE GEOPHYSICS 1/1
LAB HANSCOM AFB MA W J BURKE NOV 82

UNCLASSIFIED

END
BATT
OTTIC

OTTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

Section of the section of

Unclass SECURITY	SSIFIED CLASSIFICATI	ION AD	A1	39276	O		•	
				REPORT DOCUM		E		/ ,
1a. REPOR	T SECURITY	CLASSIFICA	ATION		16. RESTRICTIVE N			<del>/ /                                  </del>
Unclassified					10.5:55		1	
2a. SECURITY CLASSIFICATION AUTHORITY					3. DISTRIBUTION/A			
26. DECLASSIFICATION/DOWNGRADING SCHEDULE					Unlimited D	ustributio.	on .	
4. PERFORMING ORGANIZATION REPORT NUMBER(S)					5. MONITORING OF	RGANIZATION F	REPORT NUMBE	R(S)
6a. NAME C	OF PERFORMI	ING ORGAN	VIZATION	6b. OFFICE SYMBOL	7a. NAME OF MONI	TORING ORGAN	VIZATION	
	rce Geoph			(If applicable)	Air Force Ge			
	SS (City, State		de)		7b. ADDRESS (City,	State and ZIP Co.	nde)	
Hansco	m AFB, MA	1 01731			Hanscom AFB,	. MA 01731		
ORGAN	OF FUNDING/			8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT	INSTRUMENT IL	DENTIFICATION	I NUMBER
Air Force Geophysics Labora ord  Bc. ADDRESS (City, State and ZIP Code)					10. SOURCE OF FU	NDING NOS		
					PROGRAM	PROJECT	TASK	WORK UNIT
Hanscor	m AFB, MA	01731			ELEMENT NO.	NO.	NO.	NO.
	(Include Securi				7			
	atitude E		vnamics			<del></del>	<del></del>	
	nal author m J <u>. Bur</u> k							
	M J. BUTK OF REPORT	· · · · · · · · · · · · · · · · · · ·	13b. TIME C		14. DATE OF REPO		v) 15. PAGE	E COUNT
18 61.	MENTA	OTATE	FROM	то	November 19	82	one	<u> </u>
. s. surple	EMENTARY N	auon						
17.	COSATI	CODES		18. SUBJECT TERMS (C	Continue on rope-se	ecessary and id-	tify by block "	ber)
FIELD	GROUP	$\overline{}$	8. GR.		on reverse if h	, und iden	, vioce nun	
·- <del></del> -				1				
19. A8STR	ACT (Continue	; on reverse i	if necessary an	nd identify by block numbe	er)			7
							W. He	
						÷ :	•	
OTI	C FILE	: COF	γ					
20. DISTRI	BUTION/AVA	ILABILITY	OF ABSTRA	СТ	21. ABSTRACT SEC	URITY CLASSIE	ICATION	<del></del>
		_		. D DTIC USERS D		JEASSIF	<del></del>	
220. NAME OF RESPONSIBLE INDIVIDUAL					22h TELEBHONE N	22b. TELEPHONE NUMBER 22c. OFFICE SYMBOL (Include Area Code)		
			IDUAL				1	, MOUL
<u>Wil</u> lian	ı J. Burke		IDUAL				1	

## HIGH LATITUDE ELECTRODY MAMICS

William J. Burke
Air Force Geophysics
Hanscom Air Force Base, MA 01731
U.S.A.

NOS 1982

The last five years has been marked by rapid advancements in our understanding of electric fields in near-earth space. In no small part these advances have resulted from simultaneous measurements of electric fields, magnetic fields and charged particle fluxes by polar orbiting satellites above the high-latitude ionosphere. The measurement provide new evidence concerning the nature of large-scale magnetospheric convection and small scale processes leading to the field-aligned acceleration of auroral electrons. This paper reviews obervational evidence for the existence of parallel electric fields and several steady state, theoretical mechanisms by which these electric fields are sustained. Some implications of the existence of parallel electric fields for global magnetospheric modelling are also considered.

Parallel electric fields ( $E_1$ ) have occasionally been detected above the aurorae at altitudes greater than 1000 km. More often, their existence is inferred from other measurements. Intense, rapidly varying electric fields ( $\overline{E}$ ) perpendicular to the magnetic field ( $\overline{B}$ ), also known as electrostatic shocks, are frequently observed at high altitudes but not at ionospheric altitudes. Equipotential contours associated with these fields must cross magnetic field lines at some intermediate altitudes. At these altitudes equipotentials close across magnetic field lines and  $\overline{E}$  has a significant component along  $\overline{B}$ . The existence of  $E_1$  is also inferred from measurements of downcoming, auroral electron and upmoving, accelerated ion beams.

Mechanisms invoked to explain steady-state parallel electric fields include the thermoelectric effect, anomalous resistivity, quasi-neutral potentials and double layers. The theory of the last two mechanisms as well as computer simulations by Chiu and Schulz (1978) and by Wagner et al., (1981) will be discused in detail.

Field-aligned potential drops affect the moments of magnetospheric particle distributions. By opening or closing the atmospheric loss cone, E; modulates the intensity of Birkeland currents (j;).

They also affect the distribution of particle pressures in the magnetosphere. To extend magnetospheric simulation models, such as that developed by the Rice University group, to include E; requires a reexamination of the underlying physics. A variational principle will be presented from which an appropriate form of Poisson's equation NOIS (NOA) is derived and from which parallel electric fields arise in a natural principle way. The varied quantity is equivalent to the total particle and electrostatic field energy. By minimizing this quantity the particle pressure distributions that drive j; electrically coupling the ionosphere and magnetosphere, are calculated.

## References:

Chiu, Y.T. and Schulz, M.: 1978, J. Geophys. Res. 83, 629.

Wagner, J.S., Kan, J.R., Akasofu, S.-I., Tajima, T., Lebosuf, J.W. and Davson, J.M.: 1981, Physics of Auroral Arc Formation, ed. by S.-I. Akasofu and J.R. Kan, AGU, Geophysical Monograph 25, Washington, D.C. p. 304.

Climilary Avat

Avail And/o

moket

. 1 --